


Drosera ×obovata

Distribution



Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



Habitus and growth type

Height [m]: **0.05-0.2**

Life form: **hemicryptophyte**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**


Leaf arrangement (phyllotaxis): **rosulate**

Leaf shape: **simple - entire**

Stipules: **present**

Petiole: **present**

Leaf anatomy: **succulent, helomorphic**



Flower

Flowering period [month]: **June-August**

Flower colour: **white**

Flower symmetry: **actinomorphic**


Perianth type: **calyx and corolla**

Perianth fusion: **free**

Calyx fusion: **fused at the base**

Inflorescence type: **cincinnus**

Dicliny: **synoecious**



Pollination syndrome: **pseudocleistogamy**

Fruit, seed and dispersal

Fruit type: **dry fruit - capsule**

Fruit colour: **brown**

Reproduction type: **only vegetatively**



Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **native**

Habitat and sociology

Diagnostic taxon

Diagnostic taxon of associations: [RBC02 *Drosero anglicae*-*Rhynchosporium albae*](#)

Dominant taxon

Dominant taxon of associations: [RBC02 *Drosero anglicae*-*Rhynchosporium albae*](#)

Distribution and frequency

Floristic zone: **northern temperate, southern temperate, submeridional**

Floristic region: **Europe**

Elevational belt in the Czech Republic: **colline belt, submontane belt**

Occurrence frequency in the basic grid mapping cells and quadrants of the basic grid mapping cells: 16

taxon.data.freq_in_quad: 22