


Symphytum ×uplandicum

Distribution



© Petr Kouřilský

Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.



© Michal Hroneš (flora.upol.cz)



© Michal Hroneš

Habitus and growth type

Height [m]: **0.7-2**Life form: **hemicryptophyte**Life strategy: **C - competitor**

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**Stipules: **absent**Petiole: **both present and absent**Leaf life span: **summer green**Leaf anatomy: **mesomorphic, hygromorphic**

Flower

Flowering period [month]: **June-August**Flower colour: **red-violet**Perianth type: **calyx and corolla**Calyx fusion: **synsepalous**Inflorescence type: **cincinnus**Dicliny: **synoecious**Pollination syndrome: **insect-pollination, selfing**Pollinator spectrum: **bumblebees (solitary bees)**

Fruit, seed and dispersal

Reproduction type: **only by seed/spores**

Dispersal unit (diaspore): **fruit, infructescence or its part**

Dispersal strategy: **Allium (mainly autochory)**

Belowground organs and clonality

Shoot metamorphosis: **pleiocorm**

Root metamorphosis: **primary storage root**

Storage organ: **pleiocorm, primary storage root**

Shoot life span (cyclicity): **monocyclic shoots prevailing**

Branching type of stem-derived organs of clonal growth: **sympodial**

Primary root: **present**

Bud bank

Number of buds per shoot at the soil surface (root buds excluded): **5**

Number of buds per shoot at a depth of 0–10 cm (root buds excluded): **5**

Number of buds per shoot at a depth greater than 10 cm (root buds excluded): **0**

Size of the belowground bud bank (root buds excluded): **10**

Depth of the belowground bud bank (root buds excluded) [cm]: **3**

Number of buds per shoot at the soil surface (root buds included): **5**

Number of buds per shoot at a depth of 0–10 cm (root buds included): **5**

Number of buds per shoot at a depth greater than 10 cm (root buds included): **0**

Size of the belowground bud bank (root buds included): **10**

Depth of the belowground bud bank (root buds included) [cm]: **3**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Taxon origin

Origin in the Czech Republic: **neophyte**

Invasion status: **naturalized**

Geographic origin: **anecophyte**

Year of the first record in the wild: **1927**

Period of introduction: **Late Modern Period (1800-1950)**

Introduction pathway: **intentional - other**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**