



Beta vulgaris Rapacea Group

Distribution



Map info

● revised records

● unrevised records

On the map are not visualized records without the coordinates and records marked as incorrect or doubtful.

Leaf

Leaf presence and metamorphosis: **leaves present, not modified**

Petiole: **present**

Flower

Flower colour: **green**

Perianth type: **homochlamydeous**

Inflorescence type: **panicula e pseudospicis composita**



Fruit, seed and dispersal

Fruit colour: **brown**

Trophic mode

Parasitism and mycoheterotrophy: **autotrophic**

Carnivory: **non-carnivorous**

Symbiotic nitrogen fixation: **no nitrogen-fixing symbionts**

Karyology

Chromosome number (2n): **18 (27, 36)**

Ploidy level (x): **2 (3, 4)**



2C genome size [Mbp]: **1343.59**

1Cx monoploid genome size [Mbp]: **671.8**

Genomic GC content: **39.3 %**

Taxon origin

Origin in the Czech Republic: **cultivated only**

Habitat and sociology

Affinity to the forest environment

Affinity to the forest environment in Thermophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

Affinity to the forest environment in Mesophyticum and Oreophyticum: **0 - taxon that does not spontaneously occur in Czech forests**

